

Identification of a Structurally Distinct CD101 Molecule Encoded in the 950-kb *Idd10* Region of NOD Mice

Carlos Penha-Gonçalves,^{1*} Carolyn Moule,^{1*} Luc J. Smink,¹ Joanna Howson,¹ Simon Gregory,² Jane Rogers,² Paul A. Lyons,¹ Joseph J. Suttie,¹ Christopher J. Lord,¹ Laurence B. Peterson,³ John A. Todd,¹ and Linda S. Wicker¹

Genes affecting autoimmune type 1 diabetes susceptibility in the nonobese diabetic (NOD) mouse (*Idd* loci) have been mapped using a congenic strain breeding strategy. In the present study, we used a combination of BAC clone contig construction, polymorphism analysis of DNA from congenic strains, and sequence mining of the human orthologous region to generate an integrated map of the *Idd10* region on mouse chromosome 3. We found seven genes and one pseudogene in the 950-kb *Idd10* region. Although all seven genes in the interval are *Idd10* candidates, we suggest the gene encoding the EWI immunoglobulin subfamily member EWI-101 (*Cd101*) as the most likely *Idd10* candidate because of the previously reported immune-associated properties of the human CD101 molecule. Additional support for the candidacy of *Cd101* is the presence of 17 exonic single-nucleotide polymorphisms that differ between the NOD and B6 sequences, 10 causing amino acid substitutions in the predicted CD101 protein. Four of these 10 substitutions are nonconservative, 2 of which could potentially alter N-linked glycosylation. Considering our results together with those previous reports that antibodies recognizing human CD101 modulate human T-cell and dendritic cell function, there is now justification to test whether the alteration of CD101 function affects autoimmune islet destruction. *Diabetes* 52:1551–1556, 2003

From the ¹Juvenile Diabetes Research Foundation/Wellcome Trust (JDRF/WT) Diabetes and Inflammation Laboratory, Cambridge Institute for Medical Research, University of Cambridge, Addenbrooke's Hospital, Cambridge, U.K.; ²The Wellcome Trust Sanger Institute, Wellcome Trust Genome Campus, Hinxton, Cambridge, U.K.; and the ³Department of Pharmacology, Merck Research Laboratories, Rahway, New Jersey.

Address correspondence and reprint requests to Dr. Linda S Wicker, JDRF/WT Diabetes and Inflammation Laboratory, Cambridge Institute for Medical Research, Wellcome Trust/MRC Building, Addenbrooke's Hospital, Cambridge, CB2 2XY, U.K. E-mail: linda.wicker@cimr.cam.ac.uk.

Received for publication 9 August 2002 and accepted in revised form 5 March 2003.

C.J.L. is currently affiliated with the Breakthrough Breast Cancer Research Centre, Institute of Cancer Research, London, U.K.

*C.P.-G. and C.M. contributed equally to this work.

BAC, bacterial artificial chromosome; DC, dendritic cell; IEL, intraepithelial lymphocyte; FPC, fingerprint contig; MHC, major histocompatibility complex; SNP, single nucleotide polymorphism; STS, sequence-tagged site.

© 2003 by the American Diabetes Association.

The NOD mouse is increasingly recognized as an accurate and informative model of type 1 diabetes in humans. The diabetes phenotype in the NOD mouse, as found in the human disease, is under the control of numerous genetic and environmental factors. At least 17 insulin-dependent diabetes susceptibility (*Idd*) loci have been mapped in the NOD mouse through the analysis of genetic crosses involving the diabetes-resistant C57BL/6 (B6) or C57BL/10 strains (1–3). Except for the major histocompatibility complex (MHC) class II genes encoding the IA and IE molecules in *Idd1* (2) and the β 2 microglobulin locus in *Idd13* on chromosome 2 (4); no other *Idd* genes have been identified with even a moderate level of certainty.

Idd10 is located on mouse chromosome 3 and was genetically dissected from neighboring *Idd* loci using a panel of congenic strains that carry B6 donor DNA (5–8). The diabetes protective effect of the B6 allele at the *Idd10* locus in the chromosome interval as currently defined is only apparent when combined with resistance alleles at other loci. Thus, the cumulative frequency of diabetes in congenic mice bearing resistant alleles at the *Idd3* and *Idd18* loci (18% type 1 diabetes at 7 months) is decreased to 8% in combination with the *Idd10* resistance allele (8).

Previous mapping studies using NOD.B6 congenic strains defined the *Idd10* locus in the NOD genome within a 1.3-cM interval defined by the microsatellite markers *D3Mit10* and *D3Nds35* (8). Here we report on the construction and sequencing of an *Idd10* B6 bacterial artificial chromosome (BAC) clone contig and the ascertainment of its gene content. Consequently, the *Idd10* interval was more precisely defined by fine mapping the recombination break points of key congenic strains using polymorphic microsatellite markers obtained from the sequence of the region. Of the eight genes found in the *Idd10* region, the CD101 gene was selected as a candidate for *Idd10* based on its function in humans and its polymorphism in the NOD strain.

RESEARCH DESIGN AND METHODS

Congenic mouse strains. The development of the congenic mouse strains R323 (resistance allele at *Idd10*) and R93 (susceptible allele at *Idd10*) has been previously described (8).




FIG. 1. Sequence analysis of seven B6 clones in the *Idd10* region reveals seven genes and a pseudogene (bold italics). These genes are also represented in the human orthologous region 1p12. Polymorphic microsatellite markers (italics) have been identified and used to define the boundaries of the *Idd10* region using two congenic lines, susceptible R93 and resistant R323. Accession and version numbers of the B6 clones with complete sequence are as follows: RP23-335A19 (AL606750.13), RP23-356K2 (AL606757.11), RP23-267O21 (AL606744.28), RP24-357M4 (AL691436.7), RP24-369O13 (AL845553.5), RP23-357D5 (AL669872.8), and RP23-434L7 (AL645930.15).

Genotyping. Microsatellite genotyping was performed as previously described (9).

***Idd10* BAC contig construction.** To seed a BAC clone contig across the *Idd10* interval RPCI-23 library (10) (www.chori.org/bacpac), filters were hybridized with radioactively labeled probes derived from sequence-tagged sites (STSs) known to map to the region. Hybridization-positive clones were confirmed by PCR. Hybridization-positive BAC clones from the RPCI-23 library were identified within assembled fingerprint contigs (FPCs) and extracted into an *Idd10*-specific FPC database (11). The BCGSC (British Columbia Genome Sequence Centre) whole genome fingerprint database (http://www.bcgsc.bc.ca/projects/mouse_mapping/) was used as the source of initial fingerprint and contig data. Incorporation of marker data and further analyses of the fingerprint data resulted in the construction of a contiguous BAC map across the *Idd10* critical interval. Where gaps existed in the contig, end sequences of STS-positive clones were obtained from the BAC end database (www.tigr.org/tdb) and used to generate new probes to rescreen the library as above. This procedure was performed iteratively until all gaps were closed.

Genomic sequencing. To obtain the genomic sequence of the *Idd10* region, a minimal tile path of B6 BAC clones was shotgun-sequenced at the Wellcome Trust Sanger Institute.

Sequencing of mouse CD101 cDNA. RNA was extracted from unstimulated NOD spleens using RNeasy columns (Qiagen). Poly(A) RNA isolated using Genelute-mRNA miniprep kit (Sigma) was reverse transcribed (Superscript II; Invitrogen). The putative exon structure of *Cd101* was predicted using est2genome (Emboss GUI; <http://bioinfo.pbi.nrc.ca:8090/EMBOSS/index.html>) with the human *CD101* mRNA sequence NM_004258 and the B6 BAC clone RPCI-23-375D5 as input. Primers for intron-spanning PCR products were designed and the resulting PCR products sequenced (ABD). Sequence trace data were analyzed using the Staden package. Est2genome with the NOD clone DN-358O3 and the obtained cDNA sequence was used to

identify the beginning and ends of exons. The coordinates of the exons were extracted using extractseq and translated using transeq (Emboss).

Gene identification. The repeat masked B6 BAC contig sequence was subjected to blast analysis against vertebrate mRNA, dbEST, and exofish blast databases. Expressed sequence tag hits were used to identify genes within the *Idd10* region. Mouse ensembl and the orthologous human region in ensembl (www.ensembl.org/) confirmed the gene content of *Idd10*.

RESULTS

Physical delimitation of the *Idd10* genetic interval.

Physical mapping of *Idd10* was undertaken by constructing a B6 BAC clone contig covering the congenic interval previously defined as the region located between *D3Mit10* and *D3Nds35* (8). The completed minimal tile path of *Idd10* spanning and including *D3Mit10* and *D3Nds35* contained 19 BAC clones representing a size of ~3–4 Mb (data not shown). Shotgun sequence provided the opportunity to harvest potentially polymorphic microsatellites useful in refining the proximal (R323) and distal (R93) boundaries of the *Idd10* interval (Fig. 1). This iterative process narrowed the congenic interval to 950 kb and localized the *Idd10* locus distal of RP23-70A7SSR18 and proximal to *D3Mit342*. The recombination event defining the proximal boundary of *Idd10* occurs in the R323 strain and is in the 9 kb of DNA between RP23-70A7SSR18 and RP23-70A7SSR19. The distal boundary as defined by the

TABLE 1
The gene content of *Idd10* and the orthologous region on human 1p12

Human	Mouse	Description
<i>Q9NXX0</i>	<i>AK015259</i>	Ensembl predicted gene. The predicted gene is comprised of two exons. There are no domains listed, and the protein is a member of an unknown protein family.
	<i>Gapd-like</i>	Predicted by GeneWise and Genscan. The gene prediction comprises four exons. Similar to glyceraldehyde 3 phosphate.
<i>MANIA2</i>	<i>Man1b</i>	Mannosyl-oligosaccharide 1,2- α -mannosidase B. Ensembl has predicted 13 exons. The protein contains eight glycoside hydrolase domains and one coiled coil. The protein is involved in N-glycan maturation.
<i>Q9H6B2</i>	<i>FLJ22418</i>	Human ensembl predicted gene. There are seven predicted exons in human and the gene is not annotated in mouse ensembl, although FLJ22418 shows sequence similarity to the <i>Idd10</i> BAC contig. The protein contains one Ig-MHC domain, two transmembrane domains, and a signal peptide.
<i>Q9H8W5</i>	<i>FLJ13181</i>	Human ensembl predicted gene. Weakly similar to transcription intermediary factor 1 β . In human, six exons have been predicted and in mouse there are five exons predicted. The proteins contains a B-Box zinc finger, a Filamin/ABP280 repeat, an ATP/GTP-binding site motif (P-loop), and a RING finger.
<i>TTF2</i>	<i>Ttf2</i>	RNA polymerase II termination factor. Twenty-three exons have been predicted in human and only 18 exons have been predicted in mouse. The protein contains a SNF2 related domain, a helicase COOH-terminal domain, and an ATP-dependent helicase DEAH-box.
<i>CD101</i>	<i>Cd101</i>	Leukocyte surface protein. There are nine predicted exons in human and mouse. The protein contains seven Ig-MHC domains, a transmembrane domain, and a signal peptide.
<i>PTGFRN</i>	<i>Ptgfrn</i>	Prostaglandin F 2α receptor regulatory protein (FPRP). In human ensembl eight exons are predicted. The protein contains six Ig-MHC domains and a transmembrane domain. FPRP modulates the activity of the prostaglandin F 2α receptor. Found primarily in reproductive tissues, the lung, and heart.

The human gene content is taken from ensembl on 28 June 2002 (human build 29). The mouse gene list is as found in the clones sequenced in this study. All genes documented in the human analysis are supported by EST evidence. All genes in the mouse analysis are supported by EST evidence.

R93 strain is in the 82 kb between *RP23-434L7SSR1* and *D3Mit342*. Overall, the narrowing of the *Idd10* interval based on the new polymorphic markers decreased the number of BAC clones in the minimal tile path from 19 to 7.

***Idd10* gene content.** We constructed a gene map of the *Idd10* region aligning the sequence data from the seven B6 BACs in the tile path with 5.9 Mb of the human genomic sequence orthologous to *Idd10* (Fig. 1). This map provides strong evidence that the human region orthologous to *Idd10* lies in an ~900-kb region on human chromosome 1p12 (Fig. 1). Comparative analysis of sequence from the B6 tile path with the human sequence revealed eight orthologous genes within the *Idd10* interval (summarized in Table 1). Six of the seven genes with intact open reading frames within the *Idd10* interval (*Gapd* is a pseudogene) have not been reported to have particular functions in the immune system or in islet β -cells; these are *Man1b*, three genes of unknown function (*AK015259*, *FLJ22418* and *FLJ13181*), *Ttf2*, and *Fprp* (an EWI immunoglobulin subfamily member) (12,13). In contrast, *Cd101*, also an EWI subfamily member, is present on immune cells and has been partially characterized in the human immune system.

CD101 is highly expressed on monocytes, granulocytes, dendritic cells (DCs), and activated T-cells (14). In contrast to the lack of *CD101* on most resting peripheral blood lymphocytes, *CD101* is expressed on the surface of nearly all CD8+, CD3+ intestinal intraepithelial lymphocytes (IELs) (15). Interestingly, very few of these cells express CD28. Using IELs as responders, anti-*CD101* mAbs were shown to induce proliferation in conjunction with suboptimal concentrations of anti-*CD3* mAb. Therefore, *CD101* could be a costimulatory receptor functioning in this subpopulation of CD28⁻ mucosal T-cells (15). Additional studies also suggest a direct function of *CD101* on DCs (16,17). Cutaneous DCs incubated with anti-*CD101* mAb produced IL10 that led to the inhibition of T-cell proliferation (17). A direct effect of anti-*CD101* mAb on T-cell signaling has also been reported (18,19). Thus, from the gene content analysis, *CD101* is our favored candidate for further study to test its potential in mediating the effect of *Idd10* in autoimmune diabetes. However, all seven genes within the interval remain candidates until and unless it is proven that *Cd101* is *Idd10* by transgenic or knock-in approaches or other highly specific functional analyses.

NOD allotype of *CD101*. We tested for variants in the

1 atggcatgcatcctgtgtggtgcatctctctttctctctactaagttcagcatcggccagagggaaatcaagaagccctctg
1 M A C I L C V A S L F L S L T K F S I G Q R E V K I Q E G P L
94 tacagaccgaaggttacctgtcagcatcaggtgcaccgttaagtggtcatcagggctcctccacgcaggattccgggtgtctattacctg
32 Y R A E G Y P V S I R C T V S G H Q G P S T Q D F R W S I T Y L
187 ccaagcgcaccgaccaaggaagtcagatcatcagatccaaggtgcccggctctcttacgcagtgatgccagaggggtgcaagcaaggag
63 P S A P T K E V Q I I S T K D A G F S Y A V Y A Q R V Q S K E
280 atctacatagagaggctgcaggggtgactcggctcctgctcctatctcaaaactccagatgaaagatgctggcgagtacaggtgccacacacc
94 I Y I E R L Q G D S V L L H I S K L Q M K D A G E Y E C H T P
373 aacacggatgggaagtactttggaagtacagtgcaaaagcaaaatcttactgtggttcccgacaccctgtctgccaccatccctcccagagc
125 N T D G K Y F G S Y S A A K T N L T V V P D T L S A T M P S Q T
.....
466 ctcagtaagaaggaaggtgagcccttggaaactcactgtgagacaaccaagccacccgtgcaacacacccatctctctcacctgggtacctg
156 L S K K E G E P L E L T C E T T K A T V/AQ H T H L S L T W Y L
559 atgcaggaaggaggaggagccaagccactgagatcggtttctctccaaggactttgattgaccctgggtcctcctatgcagacaggtt
187 M Q E G G G S Q A T E I V S L S K D F V L T P G S S Y A D R F
652 gtggccgggtgactgacgctggacaagcttgagcaactcctcaggctgtctgtaggcaagctccagcctcagatcagggccaggtgtc
218 V A G D V R L K L G A T S F R L S V G K L Q P S D Q G Q V T
745 tgtgagccacagaatggattcaggatccagatgaaactggactttgatcacaagaaagcagacagatcaaacagctctgaggatccagccg
249 C E A T E W I Q D P D E T W T L I T R K Q T D Q T A L R I Q P
.....
838 gcagcaagagatttacagtgagcatcacagccagtagctcactgatgaaggaaaacccttggaaactggtttgctggctggtggcagagat
280 A A R D F T V S I T A S S S P D E/KG K P L E L V C L A V G R D
.....
931 ggtaacccgcagcttcaaggtgtgtggtttctcaatgggaagaaattgcccagactgatgctggtgggtcctggacctgaagagagactac
311 G N P Q L Q G V/AW F L N G K E I A Q T D A G G V L D L K R D Y
.....
1024 agagacagagcagccaagccagctgcaggtgtcaaggttaagtgccagacgttctctcaagatctctccggtgggtccagaggatgta
342 R D R A S Q G Q L Q V S K L S A Q T F S L K I F S V G P E D V
.....
1117 ggcacctacagttgtgaagtgccagaggtggcagagactcagatgggctcctggcaggtacttcagagaaagcagtcaccaggctaccgggtg
373 G T Y S C E V A E V A R T Q M G S W Q V/IL Q R K Q S P G Y R V
1210 cagctgagggagccagcaagaagtgtagccgtgtcgcagcagcagctactgtgtgggaagggagagacgctaccctctctgcaaggca
404 Q L R E P A A R S V T V S A E Q R T V W E G E T L T L L C K A
.....
1303 gctggggatgtgagtgctctatctgtgagttggtggctcacccacaggaaccagtcacacccgtgtttgtggctggatggggcaagatggt
435 A G D V S A L S V S W W L T P Q D/NO S T P V F V A G M G Q D G
1396 actgtcagctgggagttctctccgggcccctgcgaccgtggttaacaggaggctggagaaagtgactggacccttccgctggagatt
466 T V Q L G V S P G P A H R G N R R L E K V D W A T F R L E I A
1489 gcctctgccatggtcacagacagcggctacctatgaatcaggggtgtcagagagactccagaaccaggccaaggtttgcagtcaccocaaag
497 A S A M V T D S G T Y E C R V S E R L Q N Q A K G L Q S T Q K
.....
1582 atttcagtcactgtaaaatctcgaagtcaagtttacaggttaactctgatgagccgtcagccacaagtgatgctagccatccctccacctg
528 I S V T V K S L K S S L R V N L M S R Q P Q V M L A H T F H/DL
.....
1675 tctctgtagtgaggcccaactactcggatctcaagctgcggtttcggtaacgtggcagttccagccagcggctctggagccttccatcgg
559 S C V V R A N/DY S D L K L P F S V T W Q F Q P A G S G A F H R
.....
1768 ctatttcgaattgccacaatggcacogttgaatggggggagcgtcctctccagatccacaggaagcagaaggtgtcacagctcttctcttctg
590 L I R I A H N G T V E W G D V L S Q I H R K T K V S Q S F F R/H
.....
1861 tctcaactccaatctacgatcagctatggaggagacaggggtgtatcggtgtacagtgagggttatgacagagatccatagcacaagt
621 S Q L Q I Y D A A M E E T G V Y R C T V E V Y D R D S I/MC T S
.....
1954 ggcaccagcaggtatctgccacctcaacctatgaatgattactgtcaccttccagagagcaactgagcgtgaaactcaagcagtcaggtc
652 G P A R V S A T S N L L M I T V T F P E S K L S V N S S S Q V
2047 caagagctatccatcagctccagcactcagatagaatgtgccatctgtcccgatctgtggaaacctccggtatccatcatttggtacttc
683 Q E L S I S S T Q I E C A I L S R S A G N L P L S I I W Y F
2140 tcttccggtttctgcaaatgcatcctatctgaaaatctagaatggacaaaagcagtggtgtaaaatggggagcaatccaacccctcgg
714 S S V S A N A S Y L K I L E M D Q S S V V K Y G D E F Q T P R
2233 agcaagcaaaaattttctgagaaagttctcaagactattctgctgaacatctgagtggtggaggacagtgaccagggcactaccac
745 S K Q K F Y S E K V Q D L F L L N I L S V E D S D Q G H Y H
2326 tgtgctgtagaggaatggctcttctacaatgacacttggcaaaagcttgaagaaagactcaggactcacagaattgaaactcaggccc
776 C A V E E W L L S T N D T W Q K L E R K T S G L T E L K L R P
2419 acaggaagccaggtccatgtctccaaagtgaattggacaggaacgctaccgagatggagagggcgggtcagctgcagctagatggctca
807 T G S Q V H V S K V N W T G N A T E Y G E A G F S C S L D G S
.....
2512 ggcagcacagcttccctatactctgtgacatggtaccggggcagaggaactgccacggctactgctgcccagttgcaaatgccactgccacc
838 G S T A S L Y S V T W Y R G R G T A T A T A A V/TA V A N A T A T
2605 atcactgcccagcagggagtcacatctggtgacctcagctgagtgacgggtgtgctgagatggcagagaggggagcagggctccagcac
869 I T A P A G S Q M L V H L Q Y D G L L Q Y G R E G S R R L Q H
2698 tgcctaccgatctctcccacagactcgtcctgaagctgcatcgggtggaaatggaggatgctgggatatttggtgacgggtgactgagtg
900 C Y R S S P T D F V L K L H R V E M E D A G I Y W C R V T E W
2791 cagcaacatggccaccaggaagtgatcaaccaagcctcaggcagcgtcgcaacgaatggtgctcagggtgctgcgctcagagcccaggtc
931 Q Q H G H P G K W I N Q A S G E S Q R M V L R V L R S E P T V
2884 tctccctgatctgctcctcgggcccctgctccactcctcactctgctgccctctcgtcatgctgctcctctggccactgctcctctg
962 S S L I C S S G P L L H F L I V C P F V M L L L L A T S F L C
2977 ttgtaccggaagccaggaagttgtcacagctgagctcagtgcaaaagaaagaggtctctgggtgggcatgagaagacatcactcag
993 L Y R K A R K L S Q L S L S A K K E K A L W V G M R K T S L Q
3070 aaggaagctggagagagagtgacactactga
1024 K E A G E E S G H Y *

FIG. 2. The cDNA sequence of B6 *Cd101* showing the amino acid translation below. SNPs are highlighted; the NOD variants are shown above the B6 nucleotide sequence. The codon on either side of the codon affected by the polymorphism is underlined. Amino acid coding changes are highlighted, the B6 residue first followed by the NOD.

primary sequence of the CD101 protein by comparing the NOD and B6 genomic sequence. Sequencing cDNA derived from CD101 mRNA obtained from NOD spleen cells was used to establish exon/intron boundaries (because mouse CD101 had not been previously characterized) and confirm sequence variants found in the genomic analysis. Seventeen single nucleotide polymorphisms (SNPs) between NOD and B6 are found in the coding region of CD101, 10 of which resulted in amino acid changes (Fig. 2). Of these 10 changes, 4 are nonconservative and 2 of the 4 nonconservative changes could potentially alter N-linked glycosylation of the CD101 molecule by their presence in a N-X-S/T consensus sequence.

DISCUSSION

The congenic breeding strategy is a powerful method (2) to identify genes, such as *Idd10*, controlling a complex disease process exemplified in the present study by the NOD mouse model of type 1 diabetes. To reduce the number of candidate genes that must be considered, the size of the congenic region is minimized, a process that takes many years of selective breeding and phenotypic assessment. We have now reached the limit of resolution of recombination mapping that is practical for the *Idd10* interval. Considerable narrowing of the interval was gained from the sequence of the B6 BAC clones, which allowed the experimental identification of new B6/NOD polymorphic microsatellite markers. Once the informative resolving power of homologous recombination was exhausted, we were faced with the possibility of having a small physical interval that was very gene rich (20). However, only seven candidate genes were found in the *Idd10* interval (Table 1). As expected (21), nothing is known about the expression or function of half of the genes in the interval (three of the seven).

Our recent unpublished results identifying both the human type 1 diabetes locus *IDDM12* as *CTLA4* and the orthologous NOD mouse locus *Idd5.1* as *Ctla4* confirm the critical contribution of immune-related loci to the genetic control of autoimmune disease, a role that was first established by the MHC class II genes. While all genes within an interval are potential candidates, we use a triage process in which the known immune-related genes in an *Idd* interval are those that are first considered as candidates. Because definitive proof of an *Idd* gene ultimately requires a knock-in of the putative causative SNP or SNPs or a sufficiently controlled transgenic-based experiment, circumstantial evidence must be obtained for the candidacy of one of the genes within the interval and the critical nucleotide variants responsible for the disease phenotype. Hopefully, such evidence would be persuasive enough to justify the time and expense of a definitive "proof experiment."

Reports that the human CD101 molecule has costimulatory function for a subset of CD8 cells that are found primarily in the mucosal immune system of the intestine placed CD101 at the top of our *Idd10* candidate gene list. Although CD101 is essentially uncharacterized in mice, these findings in the human immune system triggered the sequencing of the CD101 gene from the NOD strain. The striking level of polymorphism found in the B6 and NOD CD101 proteins indicates it is likely that the CD101 mole-

cule is under selective pressure. These results suggest that a functional polymorphism exists for CD101 in the mouse population and that this phenotypic variation could account for the activity of *Idd10*. Sequencing of other strains of mice to determine the distribution of the CD101 variants and their haplotypic categorization is in progress. It is also important to determine the level of allelic variation of the other seven genes in the NOD *Idd10* interval by extending the NOD BAC clone shotgun sequencing or by targeted PCR product resequencing. The pattern of ancestral haplotype breakpoints in the *Idd10* region could also help to further map the causal variants (22,23). In addition, the potential association of the ortholog with human type 1 diabetes is being tested, and a knock-out of the mouse CD101 gene is underway to aid in its characterization and potentially in transgenesis experiments that may provide evidence toward the identification of *Idd10*.

CD101 is also of interest because of its potential therapeutic approachability. As has been done with other Ig superfamily members expressed at the cell surface (24,25), an engineered, soluble immunoglobulin-fusion version of CD101 designed to prevent productive engagement of CD101 or, alternatively, an agonistic antibody to augment the function of CD101 could have therapeutic effects in autoimmune disorders or in other immune-mediated processes. Pursuit of these experiments would also be an attractive way of building a case that *Cd101* might be *Idd10*.

ACKNOWLEDGMENTS

The laboratories of L.S.W., P.A.L., and J.A.T. are supported by a joint grant from the Juvenile Diabetes Research Foundation (JDRF) and the Wellcome Trust. The availability of NOD congenic mice through the Taconic Farms Emerging Models Program has been supported by grants from the Merck Genome Research Institute, National Institute of Allergy and Infectious Diseases, and the JDRF.

REFERENCES

- Lyons PA, Hancock WW, Denny P, Lord CJ, Hill NJ, Armitage N, Siegmund T, Wilson AJ, Todd JA, Phillips MS, Hess JF, Chen S-L, Fischer PA, Peterson LB, Wicker LS: The NOD *Idd9* genetic interval influences the pathogenicity of insulinitis and contains molecular variants of *Cd30*, *Tnfr2*, and *Cd137*. *Immunity* 13:107–115, 2000
- Lyons PA, Wicker LS: Localising quantitative trait loci in the NOD mouse model of type 1 diabetes. In *Genes and Genetics of Autoimmunity*. Theofilopoulos AN, Ed. Basel, Karger, 1999, p. 208–225
- Hill NJ, Lyons PA, Armitage N, Todd JA, Wicker LS, Peterson LB: The NOD *Idd5* locus controls insulinitis and diabetes and overlaps the orthologous *CTLA4/IDDM12* and *NRAMP1* loci in humans. *Diabetes* 49:1744–1747, 2000
- Hamilton-Williams EE, Serreze DV, Charlton B, Johnson EA, Marron MP, Mullbacher A, Slattery RM: Transgenic rescue implicates beta2-microglobulin as a diabetes susceptibility gene in nonobese diabetic (NOD) mice. *Proc Natl Acad Sci U S A* 98:11533–11538, 2001
- Wicker LS, Todd JA, Prins JB, Podolin PL, Renjilian RJ, Peterson LB: Resistance alleles at two non-major histocompatibility complex-linked insulin-dependent diabetes loci on chromosome 3, *Idd3* and *Idd10*, protect nonobese diabetic mice from diabetes. *J Exp Med* 180:1705–1713, 1994
- Podolin PL, Denny P, Lord CJ, Hill NJ, Todd JA, Peterson LB, Wicker LS, Lyons PA: Congenic mapping of the insulin-dependent diabetes (*Idd*) gene, *Idd10*, localizes two genes mediating the *Idd10* effect and eliminates the candidate *Fcgr1*. *J Immunol* 159:1835–1843, 1997
- Podolin PL, Denny P, Armitage N, Lord CJ, Hill NJ, Levy ER, Peterson LB, Todd JA, Wicker LS, Lyons PA: Localization of two insulin-dependent diabetes (*Idd*) genes to the *Idd10* region on mouse chromosome 3. *Mamm Genome* 9:283–286, 1998

8. Lyons PA, Armitage N, Lord CJ, Phillips MS, Todd JA, Peterson LB, Wicker LS: Mapping by genetic interaction: high-resolution congenic mapping of the type 1 diabetes loci *Idd10* and *Idd18* in the NOD mouse. *Diabetes* 50:2633–2637, 2001
9. Denny P, Lord CJ, Hill NJ, Goy JV, Levy ER, Podolin PL, Peterson LB, Wicker LS, Todd JA, Lyons PA: Mapping of the IDDM locus *Idd3* to a 0.35-cM interval containing the interleukin-2 gene. *Diabetes* 46:695–700, 1997
10. Osogawa K, Tateno M, Woon PY, Frengen E, Mammoser AG, Catanese JJ, Hayashizaki Y, de Jong PJ: Bacterial artificial chromosome libraries for mouse sequencing and functional analysis. *Genome Res* 10:116–128, 2000
11. Soderlund C, Humphray S, Dunham A, French L: Contigs built with fingerprints, markers, and FPC V4.7. *Genome Res* 10:1772–1787, 2000
12. Clark KL, Zeng Z, Langford AL, Bowen SM, Todd SC: PGRL is a major CD81-associated protein on lymphocytes and distinguishes a new family of cell surface proteins. *J Immunol* 167:5115–5121, 2001
13. Stipp CS, Kolesnikova TV, Hemler ME: EWI-2 is a major CD9 and CD81 partner and member of a novel Ig protein subfamily. *J Biol Chem* 276:40545–40554, 2001
14. Rivas A, Ruegg CL, Zeitung J, Laus R, Warnke R, Benike C, Engleman EG: V7, a novel leukocyte surface protein that participates in T cell activation. I. Tissue distribution and functional studies. *J Immunol* 154:4423–4433, 1995
15. Russell GJ, Parker CM, Sood A, Mizoguchi E, Ebert EC, Bhan AK, Brenner MB: p126 (CDw101), a costimulatory molecule preferentially expressed on mucosal T lymphocytes. *J Immunol* 157:3366–3374, 1996
16. Bagot M, Martinel I, Charue D, Weill F, Boulland ML, Wechsler J, Freeman GJ, Bensussan A, Boumsell L: CD101 is expressed by skin dendritic cells: role in T-lymphocyte activation. *Tissue Antigens* 50:439–448, 1997
17. Bouloc A, Bagot M, Delaire S, Bensussan A, Boumsell L: Triggering CD101 molecule on human cutaneous dendritic cells inhibits T cell proliferation via IL-10 production. *Eur J Immunol* 30:3132–3139, 2000
18. Soares LR, Rivas A, Tsavaler L, Engleman EG: Ligation of the V7 molecule on T cells blocks anergy induction through a CD28-independent mechanism. *J Immunol* 159:1115–1124, 1997
19. Soares LR, Tsavaler L, Rivas A, Engleman EG: V7 (CD101) ligation inhibits TCR/CD3-induced IL-2 production by blocking Ca²⁺ flux and nuclear factor of activated T cell nuclear translocation. *J Immunol* 161:209–217, 1998
20. Morel L, Blenman K, Croker B, Wakeland E: The major murine systemic lupus erythematosus susceptibility locus, *Sle1*, is a cluster of functionally related genes. *Proc Natl Acad Sci U S A* 98:1787–1792, 2001
21. Waterston RH, Lindblad-Toh K, Birney E, Rogers J, Abril JF, Agarwal P, Agarwala R, Ainscough R, Alexandersson M, An P, Antonarakis SE, Attwood J, Baertsch R, Bailey J, Barlow K, Beck S, Berry E, Birren B, Bloom T, Bork P, Botcherby M, Bray N, Brent MR, Brown DG, Brown SD, Bult C, Burton J, Butler J, Campbell RD, Carninci P, Cawley S, Chiaromonte F, Chinwalla AT, Church DM, Clamp M, Clee C, Collins FS, Cook LL, Copley RR, Coulson A, Couronne O, Cuff J, Curwen V, Cutts T, Daly M, David R, Davies J, Delehaunty KD, Deri J, Dermitzakis ET, Dewey C, Dickens NJ, Diekhans M, Dodge S, Dubchak I, Dunn DM, Eddy SR, Elnitski L, Emes RD, Eswara P, Eyas E, Felsenfeld A, Fewell GA, Flicek P, Foley K, Frankel WN, Fulton LA, Fulton RS, Furey TS, Gage D, Gibbs RA, Glusman G, Gnerre S, Goldman N, Goodstadt L, Grafham D, Graves TA, Green ED, Gregory S, Guigo R, Guyer M, Hardison RC, Haussler D, Hayashizaki Y, Hillier LW, Hinrichs A, Hlavina W, Holzer T, Hsu F, Hua A, Hubbard T, Hunt A, Jackson I, Jaffe DB, Johnson LS, Jones M, Jones TA, Joy A, Kamal M, Karlsson EK, Karolchik D, Kasprzyk A, Kawai J, Keibler E, Kells C, Kent WJ, Kirby A, Kolbe DL, Korf I, Kucherlapati RS, Kulbokas EJ, Kulp D, Landers T, Leger JP, Leonard S, Letunic I, Levine R, Li J, Li M, Lloyd C, Lucas S, Ma B, Maglott DR, Mardis ER, Matthews L, Mauceli E, Mayer JH, McCarthy M, McCombie WR, McLaren S, McLay K, McPherson JD, Meldrim J, Meredith B, Mesirov JP, Miller W, Miner TL, Mongin E, Montgomery KT, Morgan M, Mott R, Mullikin JC, Muzny DM, Nash WE, Nelson JO, Nhan MN, Nicol R, Ning Z, Nusbaum C, O'Connor MJ, Okazaki Y, Oliver K, Overton-Larty E, Pachter L, Parra G, Pepin KH, Peterson J, Pevzner P, Plumb R, Pohl CS, Poliakov A, Ponce TC, Ponting CP, Potter S, Quail M, Reymond A, Roe BA, Roskin KM, Rubin EM, Rust AG, Santos R, Sapojnikov V, Schultz B, Schultz J, Schwartz MS, Schwartz S, Scott C, Seaman S, Searle S, Sharpe T, Sheridan A, Shownkeen R, Sims S, Singer JB, Slater G, Smit A, Smith DR, Spencer B, Stabenau A, Stange-Thomann N, Sugnet C, Suyama M, Tesler G, Thompson J, Torrents D, Trevaskis E, Tromp J, Ucla C, Ureta-Vidal A, Vinson JP, Von Niederhausern AC, Wade CM, Wall M, Weber RJ, Weiss RB, Wendl MC, West AP, Wetterstrand K, Wheeler R, Whelan S, Wierzbowski J, Willey D, Williams S, Wilson RK, Winter E, Worley KC, Wyman D, Yang S, Yang SP, Zdobnov EM, Zody MC, Lander ES: Initial sequencing and comparative analysis of the mouse genome. *Nature* 420:520–562, 2002
22. Lyons PA, Armitage N, Argentina F, Denny P, Hill NJ, Lord CJ, Wilusz MB, Peterson LB, Wicker LS, Todd JA: Congenic mapping of the type 1 diabetes locus to a 780 kb region of mouse chromosome 3: Identification of a candidate segment of ancestral DNA by haplotype mapping. *Genome Res* 446–453, 2000
23. Wade CM, Kulbokas EJ 3rd, Kirby AW, Zody MC, Mullikin JC, Lander ES, Lindblad-Toh K, Daly MJ: The mosaic structure of variation in the laboratory mouse genome. *Nature* 420:574–578, 2002
24. Aruffo A, Hollenbaugh D: Therapeutic intervention with inhibitors of co-stimulatory pathways in autoimmune disease. *Curr Opin Immunol* 13:683–686, 2001
25. Egen JG, Allison JP: Cytotoxic T lymphocyte antigen-4 accumulation in the immunological synapse is regulated by TCR signal strength. *Immunity* 16:23–35, 2002